

Yellow urticaria as a rare dermatological manifestation of biliary obstruction: a case report

Mohammad Almutairi, Khalid Nabil Nagshabandi, Reem Alsergani, Sultan Alsalem

Department of Dermatology, College of Medicine, King Saud University, Riyadh, Saudi Arabia

Abstract

Urticaria is a prevalent inflammatory dermatological condition with a global estimated lifetime prevalence of up to 20%. Yellow urticaria is not a distinct medical condition; rather, it refers to the manifestation of yellow-colored wheals that appear in certain individuals with elevated serum bilirubin levels. This excess bilirubin accumulates in various tissues, including the conjunctiva and skin. The occurrence of yellow urticaria is notably rare and uncommon. In this report, we present a case of yellow-colored urticaria in a patient diagnosed with obstructive biliary stones, complemented by a concise review of previously documented cases in the literature.

Correspondence: Khalid Nabil Nagshabandi, Department of Dermatology, College of Medicine, King Saud University, PJ7C+89Q, 12372 Riyadh, Saudi Arabia.
E-mail: khaloed23@gmail.com
Tel.: +966-582940888

Key words: urticaria; yellow urticaria; bilirubin; hyperbilirubinaemia; biliary obstruction.

Conflict of interest: the authors declare no potential conflict of interest.

Ethics approval and consent to participate: ethical approval is not required for this study in accordance with local or national guidelines.

Consent for publication: written informed consent was obtained from the patient for publication of the details of his medical case and any accompanying images.

Availability of data and materials: all data generated or analyzed during this study are included in this article. Further inquiries can be directed to the corresponding author.

Received: 3 January 2025.
Accepted: 22 February 2025.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

©Copyright: the Author(s), 2025
Licensee PAGEPress, Italy
Dermatology Reports 2025; 17:10248
doi:10.4081/dr.2025.10248

Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Introduction

Urticaria is a common inflammatory skin condition with a global lifetime prevalence of up to 20%.¹ Clinically, it presents acutely with the sudden appearance of wheals, angioedema, or both, typically resolving within six weeks. In contrast, chronic urticaria is characterized by recurrent episodes of similar lesions persisting for more than six weeks.² A wheal is defined by localized superficial skin swelling, typically encircled by red erythema (flare), accompanied by itching or occasionally a burning sensation. These wheals are transient, returning to normal skin appearance within a timeframe ranging from 30 minutes to 24 hours. Urticaria may be caused by a variety of factors, usually stemming from the activation and degranulation of dermal mast cells, leading to the release of histamine and additional mediators.²

Yellow urticaria does not denote a separate disease; instead, it denotes the appearance of yellow-hued wheals observed in specific individuals with heightened levels of serum bilirubin, leading to its *surplus* accumulation in diverse tissues such as the conjunctiva and skin.³ Other factors contributing to yellow skin discoloration include carotenemia, certain medications, and inherited conditions such as Gilbert syndrome.³ The instance of yellow urticaria is uncommon and rare. Herein, we report a case of yellow-colored urticaria in a patient with obstructive biliary stones.

Case Report

A 24-year-old Saudi male with a known history of bronchial asthma presented at the emergency department at our university hospital, complaining of severe epigastric pain following a fatty meal. He reported experiencing symptoms of nausea, vomiting, pale stools, and dark urine. Physical examination revealed an unwell patient with tenderness upon palpation of the abdomen and yellow sclera. The patient's laboratory results revealed significantly elevated levels of alanine aminotransferase (ALT) at 350 U/L (normal range: 7-56 U/L) and aspartate transaminase (AST) at 185 U/L (normal range: 10-40 U/L). Alkaline phosphatase (ALP) was at 145 U/L (normal range: 44-147 U/L). Total bilirubin was markedly elevated at 116 µmol/L (normal range: 3-21 µmol/L), with direct (conjugated) bilirubin also significantly raised at 87 µmol/L (normal range: <5 µmol/L). These findings were consistent with obstructive jaundice, likely caused by biliary obstruction secondary to gallstones, as supported by the clinical presentation and imaging findings. Moreover, the patient presented with notable skin manifestations. Dermatology was consulted regarding his rash. Upon cutaneous examination, multiple urticarial lesions with a yellowish hue and edematous borders were noted over the abdomen and trunk, with positive appreciation of dermatographism (Figure 1). He also reported experiencing generalized pruritus. A skin punch biopsy was obtained for further evaluation. The management plan consisted of a 2nd-generation antihis-

tamine (loratadine) 10 mg once daily. Due to the seriousness of the patient's condition, admission to the surgical ward was necessary for further monitoring and treatment. An endoscopic retrograde cholangiopancreatography (ERCP) was performed for stone removal, aiming to relieve the obstruction in the bile duct.

Histopathological evaluation of the obtained skin lesion demonstrated an unremarkable epidermis with concomitant dermal edema along with predominantly eosinophilic infiltration within perivascular and interstitial spaces (Figure 2). Following the procedure, the yellow urticarial skin lesions disappeared entirely, nearly 24 hours after ERCP. A week post-admission, the patient developed acute pancreatitis and was transferred to the intensive care unit, where he underwent a second ERCP and was stabilized. Two weeks after the successful removal of the obstructing stones, the patient underwent an interval laparoscopic cholecystectomy to prevent recurrent biliary complications.

Discussion

This case highlights the occurrence of yellow-colored urticaria in a patient with obstructive biliary stones, shedding light on a rare and unusual manifestation of urticaria associated with elevated

serum bilirubin levels. The presentation of urticaria with yellow-hued wheals is often overlooked or misdiagnosed due to its rarity. Moreover, the association between the emergence of yellow wheals and bilirubin levels lacks clarity. While many patients displaying yellow wheals also present with hyperbilirubinemia and icteric sclera, there have been documented instances of individuals with high bilirubin levels who did not experience yellow wheals, instead exhibiting ordinary wheals without yellow discoloration.⁴

Yellow urticaria is a less frequently encountered form of urticaria, and its etiology remains elusive. It may manifest alongside hepatic and biliary disorders, with hyperbilirubinemia frequently identified, often accompanied by jaundiced sclera.² Some researchers propose that heightened serum bile acid levels might directly stimulate mast cell degranulation, initiating urticaria. Conversely, others contend that the antioxidant and anti-inflammatory properties of bilirubin could potentially mitigate urticaria symptoms.^{5,6} It is also suggested that the yellow discoloration observed in the disease may be attributed to bilirubin having an affinity for binding with elastin in the dermis, resulting in a yellowish hue similar to icteric sclera. Elastin, abundant in the sclera, typically appears yellow when bilirubin levels exceed 3 mg/dL.⁷

Antihistamines represent the primary therapeutic approach for managing yellow urticaria, mirroring the standard treatment for




Figure 1. A, B) Physical exam revealed several urticarial lesions with a yellowish hue and swollen edges across the abdomen and trunk, C) accompanied by a noticeable presence of dermatographism.




Figure 2. A) Histopathological evaluation of the skin punch biopsy revealed an unremarkable epidermis with underlying dermal edema (H&E stain, original magnification x200), and B) a higher power of the dermis shows perivascular and interstitial inflammatory cells, mainly eosinophils (H&E stain, original magnification x400).

typical non-yellowish urticaria, along with corticosteroids.² Identifying and avoiding potential triggers of urticaria is crucial for management. These triggers include physical factors such as dermographism, cold urticaria, pressure urticaria, and others, as well as dietary elements like eggs, milk, soy, and certain medications like aspirin and ACE inhibitors. Environmental factors such as infections, smoking, and pollen should also be considered. Avoiding these triggers can significantly enhance patients' quality of life.^{8,9}

This case highlights the intricate interplay between systemic conditions such as obstructive jaundice and its dermatological manifestations. The resolution of yellow urticarial lesions after ERCP suggests a direct correlation between the biliary obstruction and this dermatological manifestation. Through prompt diagnosis and appropriate intervention, our patient's symptoms were effectively managed, emphasizing the importance of a multidisciplinary approach in the care of patients with complex medical presentations.

Conclusions

Yellow-colored urticaria is a rare variant of urticaria. This case underscores the importance of recognizing and investigating this uncommon presentation of urticaria. Based on our experience, wheals of a yellowish hue serve as a clinical indicator of underlying internal disease, highlighting the necessity for healthcare practitioners to be vigilant and knowledgeable about this atypical manifestation of urticaria.

References

1. Zuberbier T, Balke M, Worm M, et al. Epidemiology of urticaria: a representative cross-sectional population survey. *Clin Exp Dermatol* 2010;35:869-73.
2. Kulthanan K, Tanwandee T, Chularojanamontri L, et al. Yellow Urticaria: A Systematic Review. *J Asthma Allergy* 2023;16:973-8.
3. Combalia A, Fustà X, Guilabert A, et al. Yellow urticaria: report of two cases and review of the literature. *J Eur Acad Dermatol Venereol* 2017;31:e204-6.
4. Lockshin NA, Hurley H. Urticaria as a sign of viral hepatitis. *Arch Dermatol* 1972;105:570-1.
5. Quist RG, Ton-Nu HT, Lillienau J, et al. Activation of mast cells by bile acids. *Gastroenterology* 1991;101:446-56.
6. Zibera L, Martelanc M, Franko M, Passamonti S. Bilirubin is an Endogenous Antioxidant in Human Vascular Endothelial Cells. *Sci Rep* 2016;6:29240.
7. Joseph A, Samant H. Jaundice. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK544252/>.
8. Alotaibi M, Khojah A, Khojah A, et al. Yellow Urticaria in a patient with gallbladder stones: A case report and review of literature. *Med Sci* 2021;25:3072-7.
9. Ackerman M, Esteve E, Potier P, Finon A. Two cases of yellow urticaria revealing acute lithiasic biliary disease. *Australas J Dermatol* 2021;62:e326-7.