

## Successful hair transplantation following repeated lipostructure in a case of morphea *en coup de sabre*: a multidisciplinary and regenerative approach

Piero Tesauro,<sup>1</sup> Paolo Tesauro,<sup>2,3</sup> Ilaria Cremonesi,<sup>1</sup> Guido Corneigliani,<sup>4</sup> Nicolò Rivetti,<sup>5</sup> Simona Muratori<sup>6</sup>

<sup>1</sup>Poliambulatorio Gioia, Milan, Italy; <sup>2</sup>Department of Otolaryngology-Head and Neck Surgery, IRCCS Azienda Ospedaliero-Universitaria of Bologna, Italy; <sup>3</sup>Department of Anesthesia, Surgery and Interventional Medicine, Gustave Roussy Institute, Villejuif, France; <sup>4</sup>Plastic Surgery Unit, MultiMedica Holding S.p.A., Reconstructive and Aesthetic Plastic Surgery School, University of Milan, Italy; <sup>5</sup>Dermatology Outpatient Clinic, Istituto Clinico Beato Matteo, Gruppo San Donato, Vigevano, Pavia, Italy; <sup>6</sup>Dermatology Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy

### Abstract

We describe the successful combination of repeated autologous lipostructure and hair transplantation in a case of morphea *en coup de sabre*, highlighting the regenerative benefits of a multidisciplinary approach. A 40-year-old woman with long-standing, stable frontoparietal and frontotemporal atrophy and alopecia underwent four fat grafting sessions using the Coleman technique over 5 years. Nine months after the final procedure, she received follicular unit extraction (FUE) hair transplantation with 1,600 grafts. The postoperative course was uneventful, with rapid and stable regrowth observed from the third month. At 12 months, the aesthetic outcome was excellent, with nearly complete graft survival and high patient satisfaction. This is the first reported case of FUE transplantation following multiple lipostructure sessions in morphea *en coup de sabre*. Fat grafting may enhance graft take by improving dermal quality and vascularization. This combined approach offers a promising solution for scarring alopecia in selected, stable patients.

**Key words:** morphea *en coup de sabre*; hair transplantation; scarring alopecia; lipostructure; follicular unit extraction.

Correspondence: Nicolò Rivetti, Dermatology Outpatient Clinic, Istituto Clinico Beato Matteo, Corso Pavia 84, 27029 Vigevano (PV), Italy.  
Tel.: +39.0381301281. E-mail: rivetti.nicolo@gmail.com

### Introduction

Morphea *en coup de sabre* is a linear variant of localized scleroderma that typically presents in childhood and progresses with periods of activity and quiescence. It affects the frontoparietal scalp and forehead, often resulting in a linear zone of alopecia, dermal atrophy, and underlying soft tissue or bony involvement. The disease is frequently disfiguring and emotionally distressing, especially in female patients.<sup>1</sup> Medical treatment during the active phase includes corticosteroids and immunosuppressants, aiming to halt progression.<sup>2,3</sup> Once the disease stabilizes, reconstructive options may be considered to address cosmetic concerns. Traditional surgical techniques, while effective, are often invasive.<sup>4-6</sup> More recently, autologous fat grafting (lipostructure) has emerged as a minimally invasive technique with both volumetric and regenerative potential.<sup>7-10</sup> Despite isolated reports of hair transplantation in this context,<sup>6,11,12</sup> no previous case has combined multiple sessions of lipostructure followed by hair restoration surgery.

### Case Report

A 40-year-old woman of Russian origin presented with two stable atrophic depressions of the scalp: a sagittal midline depression in the frontal area and a transverse depression over the right frontotemporal region (Figure 1). The first lesion appeared at 6 years of age, while the second emerged 15 years later. Neither lesion had been associated with pruritus or pain. Autoimmune screening (antinuclear antibody [ANA], extractable nuclear antigens [ENA], rheumatoid factor [RF]) was repeatedly negative. The clinical diagnosis of morphea *en coup de sabre* was made years earlier. The patient was treated with multiple courses of corticosteroids and methotrexate,<sup>2,3</sup> achieving disease stabilization. From 2018 to 2023, she underwent four autologous fat grafting procedures using the Coleman technique,<sup>10</sup> aimed at restoring soft tissue volume and improving the quality of the recipient area. Each procedure provided incremental improvement, although some degree of resorption was noted. With disease quiescence maintained for over 9 months after the final lipostructure session, surgical hair restoration was planned. The scarred recipient area measured 38 cm<sup>2</sup>, and a density

of approximately 42 follicular units per cm<sup>2</sup> was targeted. A total of 1,600 follicular units were harvested using follicular unit extraction (FUE) with a flag-pattern shave (Figure 2). Grafts were placed using Lion implanters. Due to the altered tissue consistency, graft insertion required increased precision and delicacy. Postoperative healing was uneventful. Remarkably, the patient exhibited rapid follicular regrowth beginning within 3 months, bypassing the expected telogen effluvium phase. At the 12-month follow-up, graft survival appeared close to 100%, with excellent aesthetic integration and high patient satisfaction (Figure 3).



**Figure 1.** Clinical presentation after four autologous fat grafting procedures, before hair transplantation.



**Figure 2.** Clinical presentation immediately after hair transplantation, showing both the recipient and donor areas.



**Figure 3.** Clinical presentation 12 months after hair transplantation.

## Discussion

The “*en coup de sabre*” variant of morphea poses unique challenges in aesthetic management. Linear dermal and subcutaneous atrophy, coupled with alopecia, results in a conspicuous cosmetic defect. While systemic treatments can stabilize disease activity, they offer limited benefit in reversing established scarring or alopecia.<sup>1-3</sup> Reconstructive options have traditionally included excisional surgery, skin grafts, and tissue flaps.<sup>4-6</sup> These procedures, although effective, are invasive and may not be suitable for all patients. Autologous lipostructure has gained attention for its dual volumetric and regenerative properties, particularly through the stromal vascular fraction of adipose tissue, which may modulate local inflammation and improve dermal architecture.<sup>7-10</sup> Hair transplantation in scarring alopecia remains controversial, with graft survival heavily dependent on recipient bed quality. Literature on hair transplantation in morphea *en coup de sabre* is limited to three cases.<sup>11-13</sup> Park *et al.*<sup>11</sup> reported 85% graft survival at 10 months. Dai *et al.*<sup>12</sup> reported 86.7% at 12 months. A third case initially showed positive results, but graft failure occurred at 4-year follow-up (personal communication). To our knowledge, our case is the first to combine multiple lipostructure sessions with subsequent FUE transplantation. The regenerative effect of fat grafting likely enhanced the vascularity and elasticity of the scarred tissue, improving graft retention. Rapid follicular regrowth and long-term stability suggest this multidisciplinary approach may offer superior outcomes in selected patients.

## Conclusions

Hair transplantation in stabilized morphea *en coup de sabre* is feasible and can yield excellent cosmetic results. Pretreatment with repeated autologous fat grafting may significantly enhance the recipient site, facilitating graft survival and natural integration. This case underscores the value of combining regenerative and surgical strategies in managing cicatricial alopecia. Further studies are warranted to validate these findings and establish standardized protocols.

## References

1. Kreuter A, Moinzadeh P, Kinberger M, et al. S2k guideline: Diagnosis and therapy of localized scleroderma. *J Dtsch Dermatol Ges* 2024;22:605-20.
2. Weibel L, Sampaio MC, Visentin MT, et al. Evaluation of methotrexate and corticosteroids for the treatment of localized scleroderma (morphoea) in children. *Br J Dermatol* 2006;155:1013-20.
3. Zulian F, Martini G, Vallongo C, et al. Methotrexate treatment in juvenile localized scleroderma: a randomized, double-blind, placebo-controlled trial. *Arthritis Rheum* 2011;63:1998-2006.
4. Ayoub R, Saba SC. Treatment of linear scleroderma "En coup de Sabre" with single-stage autologous fat grafting: a case report and review of the literature. *J Cosmet Dermatol* 2021;20:285-9.
5. El Omari M, Debbarh M, Lakhdari MA, et al. Adipose tissue grafting for the treatment of morphea en coup de sabre: a simple filler or an emerging cellular therapy? *Cureus* 2022;14:e30358.
6. Pierre-Louis M, Sperling LC, Wilke MS, Hordinsky MK.

- Distinctive histopathologic findings in linear morphea (en coup de sabre) alopecia. *J Cutan Pathol* 2013;40:580-4.
7. Klinger M, Marazzi M, Vigo D, Torre M. Fat injection for cases of severe burn outcomes: a new perspective of scar remodeling and reduction. *Aesthetic Plast Surg* 2008;32:465-9.
  8. Tremolada C, Colombo V, Ventura C. Adipose tissue and mesenchymal stem cells: state of the art and Lipogems® technology development. *Curr Stem Cell Rep* 2016;2:304-12.
  9. Debuc B, Gendron N, Cras A, et al. Improving autologous fat grafting in regenerative surgery through stem cell-assisted lipotransfer. *Stem Cell Rev Rep* 2023;19:1726-54.
  10. Coleman SR. Structural fat grafting. *Aesthet Surg J* 1998;18:386-8.
  11. Park SW, Wang HY. Survival of grafts in coup de sabre. *Dermatol Surg* 2002;28:763-6.
  12. Dai Y, Xu AE, He J. A case report of coup de sabre patient with hair transplantation. *J Dermatolog Treat* 2017;28:546-8.
  13. Ekelem C, Pham C, Atanaskova Mesinkovska N. A systematic review of the outcome of hair transplantation in primary scarring alopecia. *Skin Appendage Disord* 2019;5:65-71.

---

Received: 4 May 2025; Accepted: 11 May 2025.

Contributions: Piero Tesauro: hair transplantation procedure; Paolo Tesauro, Nicolò Rivetti, Ilaria Cremonesi: follow-up and manuscript preparation; Guido Cornegliani: lipostructure sessions; Simona Muratori: initial management and prescription of systemic therapies. All authors have made substantial intellectual contributions to this work, have reviewed and approved the final version of the manuscript, and agree to be accountable for all aspects of the study.

Conflict of interest: the authors have no conflict of interest to declare.

Ethics approval and consent to participate: this case report did not require formal ethical committee approval. Written informed consent was obtained from the patient for all diagnostic and therapeutic procedures.

Consent for publication: the patient gave her written consent to use her personal data for the publication of this case report and any accompanying images.

Availability of data and materials: all data generated or analyzed during this study are included in this published article.

Acknowledgments: we thank the patient for consenting to the publication of this case and the accompanying photographic documentation.

*Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

*This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).*