

Pilot study of topical management of folliculitis: clinical outcomes and safety in routine practice

Meda-Elena Stefancu,¹ Dionisio Franco Barattini,² Ionel Botnaru,³ Carmen Vizman,⁴ Luca Stucchi,⁵
Luca Barattini⁶

¹SC Salvosan Ciobanca S.r.l., Zalău, Romania; ²Opera CRO (Tigermed EMEA), Timișoara, Romania;

³Darzas Aesthetic, Oradea, Romania; ⁴Nova-Clin Medical Research Center S.r.l., Timișoara, Romania;

⁵BMG Pharma S.p.A., Milan, Italy; ⁶Tigermed Italy S.r.l., IT Department, Genoa, Italy

doi: 10.4081/dr.2026.10810

SUPPLEMENTARY MATERIAL

Supplementary Table 1. Change in the DLQI total score at 8 weeks after the initiation of treatment compared to baseline.

Does folliculitis affect the patient's life?	Day 0 visit N (%)	Day 56±4 visit N (%)
No effect	1 (7.7)	7 (53.8)
Small effect	3 (23.1)	5 (38.5)
Moderate effect	4 (30.8)	0 (0.0)
Large effect	0 (0.0)	0 (0.0)
Very large effect	4 (30.8)	1 (7.7)
Extremely large effect	1 (7.7)	0 (0.0)

p<0.001 between Day 0 and Day 56±4 (Bhapkar test).